

CLAIMS

What is claimed is:

1. A method for identifying inherited point mutations in a target region of a genome, comprising providing a pool of DNA fragments isolated from a population, and
 - a) amplifying said target region of each of said fragments in a high fidelity polymerase chain reaction (PCR) under conditions suitable to produce double stranded DNA products which contain a terminal high temperature isomelting domain that is labeled with a detectable label, and where the mutant fraction of each PCR-induced mutation is not greater than about 5×10^{-5} ;
 - b) melting and reannealing the product of a) under conditions suitable to form duplexed DNA, thereby producing a mixture of wild type homoduplexes and heteroduplexes which contain point mutations;
 - c) separating the heteroduplexes from the homoduplexes based upon the differential melting temperatures of said heteroduplexes and said homoduplexes and recovering the heteroduplexes, thereby producing a second pool of DNA that is enriched in target regions containing point mutations;
 - d) amplifying said second pool in a high fidelity PCR under conditions where only homoduplexed double stranded DNA is produced, thereby producing a mixture of homoduplexed DNA containing wild type target region and homoduplexed DNAs which contain target regions that include point mutations;
 - e) resolving the homoduplexed DNAs containing target regions which include point mutations based upon the differential melting temperatures

-171-

of the DNAs, and recovering the resolved DNAs which contain a target region which includes point mutations; and

- f) sequencing the target region of the recovered DNAs to identify point mutations within the target region.

5 2. The method of Claim 1 wherein said population comprises at least 1000 individuals.

3. The method of Claim 1 wherein said population comprises at least 10,000 individuals.

10 4. The method of Claim 1 wherein said population comprises between about 10,000 and about 1,000,000 individuals.

5. The method of Claim 1 wherein said population is a population of humans.

6. The method of Claim 5 wherein said human population consists of members of the same demographic group.

15 7. The method of Claim 6 wherein said human population consists of individuals of European ancestry or a subgroup thereof.

8. The method of Claim 6 wherein said human population consists of individuals of African ancestry or a subgroup thereof.

9. The method of Claim 6 wherein said human population consists of individuals of Asian ancestry or a subgroup thereof.

-172-

10. The method of Claim 6 wherein said human population consists of individuals of Indian ancestry or a subgroup thereof.
11. The method of Claim 1 wherein said pool is enriched in fragments containing said target region.
- 5 12. The method of Claim 1 wherein said high fidelity PCR is catalyzed by a high fidelity polymerase, and wherein the copy number of each target region in a) is doubled a maximum of about 6 times.
- 10 13. The method of Claim 1 wherein the heteroduplexes are separated from the homoduplexes in c), and the homoduplexed DNAs are resolved in d) by constant denaturing gel capillary electrophoresis, constant denaturing gel electrophoresis, denaturing gradient gel electrophoresis or denaturing high performance liquid chromatography.
- 15 14. The method of Claim 1 wherein the heteroduplexes are separated from the homoduplexes in c), and the homoduplexed DNAs are resolved in d) by constant denaturing gel capillary electrophoresis.
15. The method of Claim 1 wherein the target region is an isomelting domain consisting of about 80 to about 3,000 base pairs (bp).
16. The method of Claim 1 wherein the target region is about 80 to about 1000 bp.
17. The method of Claim 1 wherein the target region is about 100 to about 500 bp.
- 20 18. The method of Claim 1 wherein the target region an exon or portion thereof of a protein encoding gene.

19. The method of Claim 1 wherein the target region spans the junction of an intron and an exon of a protein encoding gene.
20. The method of Claim 1 wherein the target region is a regulatory region of a protein encoding gene.
- 5 21. The method of Claim 1 wherein the target region is a intron of a protein encoding gene.
22. The method of Claim 1 wherein said target region is in an gene which encodes RNA.

2025 RELEASE UNDER E.O. 14176

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

23. A method for identifying genes which carry a harmful allele, comprising:
- identifying the inherited point mutations which are found in the genes or portions thereof of a population of young individuals, determining the frequencies with which each point mutation occurs, and calculating the sum of the frequency of all point mutations identified for each gene or segment;
 - identifying the inherited point mutations which are found in the genes or portions thereof of a population of aged individuals, determining the frequencies with which each point mutation occurs, and calculating the sum of the frequencies of all point mutations identified for each gene or segment;
 - comparing the sum frequency of point mutations which are found in a selected gene or portion thereof of the young population calculated in a) with the sum frequency of point mutation which are found in the same gene or portion thereof of the aged population calculated in b), wherein a significant decrease in the sum frequency of point mutations in the aged population indicates that said selected gene carries a harmful allele.
24. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 23.

25.

A method for identifying genes which carry a harmful allele, comprising:

- a) identifying the inherited point mutations which are found in the genes or portions thereof of a population of young individuals, and determining the frequencies with which each point mutation occurs;
- b) identifying the inherited point mutations which are found in the genes or portions thereof of a population of aged individuals, and determining the frequency with which each point mutation occurs; and
- c) comparing the frequency of each point mutation identified in a selected gene or portion thereof of the young population determined in a) with the frequency of the same point mutations identified in said selected gene of the aged population determined in b), wherein a significant decrease in the frequency of two or more point mutations in said selected gene of the aged population relative to said selected gene of the young population indicates that said selected gene carries a harmful allele.

15

26.

The method of Claim 25 further comprising:

- d) determining the frequency of said two or more point mutations which decrease in the aged population in said selected gene of one or more intermediate age-specific populations;
- e) determining the age-specific decline of said two or more point mutations; and
- f) comparing the age-specific decline determined in e) with the theoretical age-specific decline of harmful alleles which cause mortal diseases, $X(h,t)$, and determining if the functions are significantly different, wherein a determination that the age-specific decline determined in e) is not significantly different from the theoretical age-specific decline of harmful alleles which cause one or more mortal diseases further indicates that said selected gene carries a harmful allele and has a high probability of being causal of said one or more mortal diseases.

27. The method of Claim 26 further comprising:

- g) determining the frequency of said two or more point mutations which decrease in the aged population in said selected gene of one or more proband populations; and
- h) comparing the frequencies of said two or more point mutations in said selected gene or portion thereof in the young population with the frequencies of said two or more point mutations in said selected gene or portion thereof in the proband populations; wherein a significant increase in the frequencies of said one or more point mutations in the proband population relative to the young population indicates that said gene carries a harmful allele that plays a causal role in said disease.

28. The method of Claim 26 further comprising:

- g) determining the frequency of said two or more point mutations which decrease in the aged population in said selected gene of one or more proband populations consisting of individuals with early onset disease; and
- h) comparing the frequencies of said two or more point mutations in said selected gene or portion thereof in the young population with the frequencies of said two or more point mutations in said selected gene or portion thereof in the proband populations; wherein a significant increase in the frequencies of said one or more point mutations in the proband population relative to the young population indicates that said gene carries a harmful allele which is a secondary risk factor which accelerates the appearance of disease.

29. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 25.

- [illegible]

33. A method for identifying genes which carry a harmful allele or which are linked to a gene that carries a harmful allele, comprising:
- a) identifying the inherited point mutations which are found in the genes or portions thereof of a population of young individuals, and determining the frequency with which each point mutation occurs;
 - b) identifying the inherited point mutations which are found in the genes or portions thereof of a population of aged individuals, and determining the frequency with which each point mutation occurs;
 - c) comparing the frequency of each point mutation identified in a selected gene or portion thereof of the young population determined in a) with the frequency of the same point mutations identified in said selected gene of the aged population determined in b), wherein a significant decrease in the frequency of a point mutation in said selected gene of the aged population relative to said selected gene of the young population indicates that said selected gene carries a harmful allele or is linked to a gene that carries a harmful allele.
34. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 33.
35. A method of identifying genes which carry a harmful allele that is a secondary risk factor that accelerates the appearance of a disease, comprising:
- a) identifying the inherited point mutations which are found in the genes or portions thereof of an early onset proband population, and determining the frequency with which each point mutation occurs;
 - b) identifying the inherited point mutations which are found in the genes or portions thereof of a late onset proband population, and determining the frequency with which each point mutation occurs;
 - c) comparing the frequencies of point mutations which are found in a

selected gene or portion thereof in the early onset proband population with the frequencies of the same point mutations in said selected gene or portion thereof of the late onset proband populations; wherein a significant increase in the frequencies of one or more point mutations in the early onset proband population relative to the late onset proband population indicates that said gene carries a harmful allele which is a secondary risk factor which accelerates the appearance of disease.

5

36. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 35.

10

37. A method of identifying genes which carries a harmful allele that is a secondary risk factor that accelerates the appearance of a disease, comprising:

15

a) identifying the inherited point mutations which are found in the genes or portions thereof of an early onset proband population, determining the frequency with which each point mutation occurs, and calculating the sum of the frequency of all point mutations identified for each gene or segment;

20

b) identifying the inherited point mutations which are found in the genes or portions thereof of a late onset proband population, and determining the frequency with which each point mutation occurs, and calculating the sum of the frequency of all point mutations identified for each gene or segment;

25

c) comparing the sum frequency of point mutations which are found in a selected gene or portion thereof of the early onset proband population calculated in a) with the sum frequency of point mutation which are found in the same gene or portion thereof of the late onset proband population calculated in b), wherein a significant decrease in the sum frequency of point mutations in the late onset proband population

-180-

indicates that said selected gene carries a harmful allele which is a secondary risk factor that accelerates the appearance of a disease.

38. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim.37.
- 5 39. A method for identifying genes which carry an allele which increases longevity, comprising:
- 10 a) identifying the inherited point mutations which are found in the genes or portions thereof of a population of young individuals, determining the frequencies with which each point mutation occurs, and calculating the sum of the frequency of all point mutations identified for each gene or segment;
- 15 b) identifying the inherited point mutations which are found in the genes or portions thereof of a population of aged individuals, determining the frequencies with which each point mutation occurs, and calculating the sum of the frequencies of all point mutations identified for each gene or segment;
- 20 c) comparing the sum frequency of point mutations which are found in a selected gene or portion thereof of the young population calculated in a) with the sum frequency of point mutation which are found in the same gene or portion thereof of the aged population calculated in b), wherein a significant increase in the sum frequency of point mutations in the aged population indicates that said selected gene carries an allele which increases longevity.
40. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 39.
- 25

41. A method for identifying genes which carry an allele which increases longevity, comprising:
- a) identifying the inherited point mutations which are found in the genes or portions thereof of a population of young individuals, and determining the frequencies with which each point mutation occurs;
 - b) identifying the inherited point mutations which are found in the genes or portions thereof of a population of aged individuals, and determining the frequency with which each point mutation occurs; and
 - c) comparing the frequency of each point mutation identified in a selected gene or portion thereof of the young population determined in a) with the frequency of the same point mutations identified in said selected gene of the aged population determined in b), wherein a significant increase in the frequency of two or more point mutations in said selected gene of the aged population relative to said selected gene of the young population indicates that said selected gene carries an allele which increases longevity.
42. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 41.

43. A method for identifying genes which carry an allele which increases longevity or which are linked to a gene that increases longevity, comprising:
- a) identifying the inherited point mutations which are found in the genes or portions thereof of a population of young individuals, and determining the frequency with which each point mutation occurs;
 - b) identifying the inherited point mutations which are found in the genes or portions thereof of a population of aged individuals, and determining the frequency with which each point mutation occurs;
 - c) comparing the frequency of each point mutation identified in a selected gene or portion thereof of the young population determined in a) with the frequency of the same point mutations identified in said selected gene of the aged population determined in b), wherein a significant increase in the frequency of a point mutation in said selected gene of the aged population relative to said selected gene of the young population indicates that said selected gene carries an allele which increases longevity or is linked to a gene that increases longevity.
44. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 43.

45. A method for identifying genes which affect the incidence of a disease, comprising:
- 5 a) identifying the inherited point mutations which are found in genes or portions thereof of a population of young individuals not afflicted with said disease, determining the frequencies with which each point mutation occurs, and summing the frequency of all point mutations identified in each gene or segment thereof;
- 10 b) identifying the inherited point mutations which are found in genes or portions thereof of a proband population having said disease, determining the frequencies with which each point mutation occurs, and summing the frequency of all point mutations identified in each gene or segment thereof;
- 15 c) comparing the sum frequency of point mutation in a selected gene or portion thereof in the young population with the sum frequency of point mutations in said selected gene or portion thereof in the proband population; wherein a significant increase in the sum frequency of point mutations in the proband population indicates that said gene plays a causal role in said disease.
46. The method of Claim 29 wherein said disease is a mortal disease.
- 20 47. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 45.

48. A method for identifying a gene which carries deleterious alleles, comprising
- a) identifying the inherited point mutations occurring in the exon(s) and splice sites of said gene of a population of young individuals;
 - b) identifying the subset of point mutations in a) that are obligatory knockout point mutations, and determining the frequencies with which each obligatory knockout point mutation occurs; and
 - c) summing the frequency of all obligatory knockout point mutations identified in the gene; wherein a sum frequency of less than about 2% indicates that said gene carries a deleterious allele.
- 10 49. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 48.
50. A method for identifying a gene which carries deleterious alleles, comprising
- a) identifying the inherited point mutations occurring in the exon(s) and splice sites of said gene of a population of young individuals;
 - 15 b) identifying the subset of point mutations in a) that are obligatory knockout point mutations, and determining the frequencies with which each obligatory knockout point mutation occurs;
 - c) identifying the subset of point mutations in a) that are presumptive knockout point mutations, and determining the frequencies with which each presumptive knockout point mutation occurs; and
 - 20 d) summing the frequency of all of said obligatory knockout point mutations and presumptive knockout point mutations identified in the gene; wherein a sum frequency of less than about 2% indicates that said gene carries a deleterious allele.
- 25 51. The method of Claim 32 wherein a sum of about 0.02% to about 2% indicates that said gene carries a recessive deleterious allele.

-185-

52. The method of Claim 32 wherein a sum of less than about 0.02% indicates that said gene carries a dominant deleterious allele.
53. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 50.

00411000-004

54. A method for isolating and identifying a target region of a genome which contains inherited point mutations, comprising providing a pool of DNA fragments isolated from a population, and
- 5 a) amplifying said target region of each of said fragments in a high fidelity polymerase chain reaction (PCR) under conditions suitable to produce double stranded DNA products which contain a terminal high temperature isomelting domain that is labeled with a detectable label, and where the mutant fraction of each PCR-induced mutation is not greater than about 5×10^{-5} ;
 - 10 b) melting and reannealing the product of a) under conditions suitable to form duplexed DNA, thereby producing a mixture of wild type homoduplexes and heteroduplexes which contain point mutations;
 - c) separating the heteroduplexes from the homoduplexes based upon the differential melting temperatures of said heteroduplexes and said
15 homoduplexes and recovering the heteroduplexes, thereby producing a second pool of DNA that is enriched in target regions containing point mutations;
 - d) amplifying said second pool in a high fidelity PCR under conditions where only homoduplexed double stranded DNA is produced, thereby
20 producing a mixture of homoduplexed DNA containing wild type target region and homoduplexed DNAs which contain target regions that include point mutations;
 - e) resolving the homoduplexed DNAs containing target regions which include point mutations based upon the differential melting temperatures of the DNAs, and recovering the resolved DNAs which contain a target
25 region which includes point mutations; and
 - f) sequencing the target region of a recovered DNA which contain a target region which include point mutations.

-187-

55. An isolated target region of a genome which contains an inherited point mutations isolated by the method of Claim 54.
56. An isolated nucleic acid which is complimentary to a strand of a gene or allele thereof identified by the method of Claim 54.
- 5 57. An array of isolated nucleic acids, immobilized on a solid support, said array having at least about 100 different isolated nucleic acids which occupy separate known sites in said array, wherein each of said different isolated nucleic acids hybridizes to a target region which contains an inherited point mutation of Claim 54.
- 10 58. An array of isolated nucleic acids, immobilized on a solid support, said array having at least about 100 different isolated nucleic acids which occupy separate known sites in said array, wherein said array comprises all known deleterious, harmful and beneficial point mutations for all human populations.

59. A method of identifying the inherited point mutations in any target region of a genome of a population, wherein said point mutations
- a) interfere with reproduction;
 - b) cause or accelerate the appearance of a mortal disease; or
 - c) prevent or delay the appearance of a mortal disease; wherein
- the set of all inherited point mutations occurring at a frequency at or above 5×10^{-5} is first identified separately in members of the same population who comprise subpopulations selected from the group consisting of young, aged, intermediate age, afflicted with disease, afflicted with a disease of early age onset and afflicted with a disease of late age onset, by noting the frequencies of each inherited point mutation within and between the subpopulations, thereby determining which inherited point mutations are deleterious, harmful or beneficial.

add
B2

ADD
D6

SUB
DB

2025-10-10 10:10:10